

Multi-particle States in Spin-1 Chain System CsNiCl₃

M. Kenzelmann¹, R. A. Cowley¹, W. J. L. Buyers^{2,3}, R. Coldea^{4,5}, J. S. Gardner², M. Enderle⁶, D. F. McMorrow⁷, S. M. Bennington⁵

(1) Oxford Physics, Clarendon Laboratory, Oxford OX1 3PU, UK (2) Neutron Program for Materials Research, National Research Council of Canada, Chalk River, Ontario, Canada K0J 1J0 (3) Canadian Institute for Advanced Research (4) Oak Ridge National Laboratory, Solid State Division, Oak Ridge, TN 37831, USA (5) ISIS Facility, Rutherford Appleton Laboratory, Oxon OX11 0QX, UK (6) Technische Physik, Geb. 38, Universität des Saarlandes, 66123 Saarbrücken, Germany (7) Condensed Matter Physics and Chemistry Department, Risø National Laboratory, DK-4000, Roskilde, Denmark
(February 1, 2008)

A continuum of magnetic states has been observed by neutron scattering from the spin-1 chain compound CsNiCl₃ in its disordered gapped one-dimensional phase. Results using both triple-axis and time-of-flight spectrometers show that around the antiferromagnetic point $Q_c = \pi$, the continuum lies higher in energy than the Haldane gapped excitations. At 6 K the integrated intensity of the continuum is about 12(2)% of the total spectral weight. This result is considerably larger than the 1-3% weight predicted by the non-linear sigma model for the 3-particle continuum.

PACS numbers: 75.25.+z, 75.10.Jm, 75.40.Gb

The excitations of one-dimensional (1D) Heisenberg antiferromagnets have attracted much experimental and theoretical attention ever since Haldane [1] predicted that the excitations of integer spin and half-integer spin chains are different. Half-integer spin chains have no spin gap and exhibit a spinon continuum extending to zero energy [2,3]. For integer-spin chains the spectrum is at low temperatures dominated by well-defined single particle excitations corresponding to a triplet of spin-1 particles that exhibit a large energy gap [4,5]. These have been thoroughly studied by neutron scattering from spin-1 chain compounds and it has been shown that the well-defined excitations largely exhaust the total scattering [6]. Nonetheless, a continuum of multi-particle scattering was predicted by Haldane and others [7], but as yet no experiment has established its existence. To search for such a continuum we have measured the inelastic neutron scattering of CsNiCl₃ in its paramagnetic 1D phase using time-of-flight and triple-axis spectrometers. We find a neutron scattering continuum at energies higher than the well-defined single particle excitation, whose integrated intensity is bigger than predicted for the multi-particle scattering.

CsNiCl₃ is magnetically a quasi 1D spin-1 chain compound with the Ni chains along the c -axis of the hexagonal unit cell. Each unit cell contains two Ni atoms and the Hamiltonian is

$$H = J \sum_i^{\text{chain}} \vec{S}_i \cdot \vec{S}_{i+1} + J' \sum_{\langle i,j \rangle}^{\text{plane}} \vec{S}_i \cdot \vec{S}_j - D \sum_i (\vec{S}_i^z)^2. \quad (1)$$

The exchange interaction $J = 2.28$ meV along the c -axis is much stronger than the exchange interaction in the basal plane $J' = 0.044$ meV [4,8,9]. The weak Ising anisotropy $D = 4$ μ eV is small enough that CsNiCl₃ is a good example of an isotropic Heisenberg antiferromagnet. Below $T_N = 4.85$ K the interchain coupling causes long-range ordering of the magnetic moments. Above T_N

CsNiCl₃ is in a 1D magnetic phase for which the magnetic exchange interaction along the c -axis dominates the spin dynamics, except close to T_N .

The sample of CsNiCl₃ was a single crystal 20 mm $\times$ 5 mm $\times$ 5 mm and was mounted with its (hhl) crystallographic plane in the principal scattering plane. Inelastic neutron scattering experiments were carried out with the time-of-flight spectrometer, MARI, at the ISIS facility of the Rutherford Appleton Laboratory and with reactor-based triple-axis spectrometers, DUALSPEC at the Chalk River Laboratories and RITA at Risø National Laboratory.

For the experiment using MARI the incoming neutron energy was 20 and 30 meV. The energy resolution was 0.35 and 0.4 meV, as determined from the full width at half maximum (FWHM) of the quasi-elastic peak. The resolution in wave-vector transfer at zero energy transfer was typically 0.02 $\text{\AA}^{-1}$ along the c^* -axis and along the [110] direction and up to 0.19 $\text{\AA}^{-1}$ perpendicular to the scattering plane if only the central detector bank was used. Both the energy and the wave-vector resolution improved with increasing energy transfer. The measurements were performed with the sample at temperatures between 6.2 and 12 K.

The measurements at the DUALSPEC triple-axis spectrometer were performed at 8.5 K with a fixed scattered neutron energy of 14.51 meV and with a graphite filter to absorb the higher order reflections from the pyrolytic graphite monochromator and analyzer. The collimation from reactor to detector was 0.65°-0.6°-1.4°-2.0° and gave an energy resolution of 1.94 meV, as determined from the quasi-elastic peak. In the case of the RITA triple-axis spectrometer, supermirror guides and a rotating velocity selector produced a variable energy monochromatic neutron beam with suppressed higher order contamination. The scattered neutrons were analyzed using the Soller geometry with a fixed energy of 5 meV and a cooled beryllium filter between the sam-

ple and analyzer. The effective collimation was 1.45° – 1.45° – 1° – 2.0° and the energy resolution was 0.35 meV as determined from the quasi-elastic peak.

The measurements with MARI at temperatures between 6.2 and 12 K were performed with the c-axis of the sample either perpendicular or parallel to the incoming beam. The dynamic structure factor $S(\vec{Q}, E)$ was measured for wave-vector transfers along the c-axis between $l \simeq 0.175$ and ~ 1.5 , where the wave-vector along the chain is $\frac{2\pi l}{c}$. Because there are two Ni atoms in each unit cell along c, $l = 1$ corresponds to the antiferromagnetic (AF) wave-vector, π , generally used in theoretical work on AF chains. The data with $l > 1.5$ was contaminated by phonon scattering and so no analysis of the magnetic contribution was then attempted.

FIG. 1. The neutron scattering intensity observed using MARI as a function of energy transfer and wave-vector transfer along the [001] direction. The data shown is an average of measurements at 6.2 and 12 K and is given on a logarithmic scale. The intense band of scattering arises from the well-defined Haldane modes and the scattering continuum is at higher energies up to ~ 12 meV for $0.6 < l < 1.4$. The solid lines are the boundary of the continuum that would be observed for an $S=1/2$ linear chain if the maximum of the lower boundary was at the same energy as the Haldane excitation for $l = 0.5$.

Constant- l scans were constructed from the measured $S(\vec{Q}, E)$ by mapping on (l, E) -space (Fig. 1). This data was then averaged in stripes with a width $\Delta l = 0.05$ – 0.10 and corrected for neutron absorption. Fig. 1 shows directly the continuum scattering. It is stronger at AF momenta around $l = 1$ than in the region near $l = 0.2$, which is known to carry vanishing magnetic weight [11], and the zone boundary region $l = 0.5$. A detailed description of the analysis of the experiment will be given elsewhere [10]. Here, we focus only on the scattering close to the AF point as measured in the central detector bank. As shown in Fig. 2a) the response at $Q_c = \pi$ consists of the sharp mode observed previously but also has a component that extends to high frequencies. This continuous spectrum has not been observed before. It is well above background and moreover is seen to be much larger than the scattering near $Q_c = 0$ where the magnetic scattering is expected [11] to be vanishingly small. We will show below that the continuum cannot arise from resolution broadening of the well-defined mode.

The magnetic excitations in the MARI experiment were best fitted by an antisymmetrized Lorentzian peak weighted with the detailed balance factor and convoluted with the line-shape of the quasi-elastic incoherent scattering, which is a good estimate of the resolution line-shape particularly at low energy transfers. An intrinsic Lorentzian peak is supported by a recent theoretical prediction [12] and a recent experiment [13]. It gives an excellent fit to the low energy resonant Haldane peak.

Nevertheless we see in Fig. 2a) that the scattering intensity at energies above the gap energy is significantly higher than expected from the fits to the sharp peak. The continuum scattering is clearly observed for $0.6 < l < 1.4$ (Fig. 1). At 6.2 K, the integrated intensity of the continuum is 10(4)% of the total magnetic scattering at the AF point. The continuum was also observed in an experiment in a different configuration at 12 K, where the c-axis was parallel to the incoming beam.

Calculations were made using an ISIS spectrometer simulation programme which takes into account the spectrometer parameters such as the detailed pulse line-shape, chopper characteristics etc. and predicts the scattering line-shape for a particular sample orientation and model of the scattering cross-section. We find that the resolution function reproduced the measured line-shape of the quasi-elastic incoherent scattering and of the well-defined excitations, but that it cannot give rise to a high-energy tail in the region of the continuum.

FIG. 2. a) Neutron scattering intensity at 6.2 K as a function of energy transfer at $l = 0$ and $l = 1$ as measured using the time-of-flight spectrometer MARI at ISIS and after correction for neutron absorption. The wave-vector transfer describing the in-plane wave-vector [hh0] varies between $h = 0.09$ and 0.81 . The solid line is the fit described in the text. b) Neutron scattering counts at 8.5 K as a function of energy transfer as measured using the triple-axis spectrometer DUALSPEC and corrected for neutron absorption. The energy scan at the zone boundary (0.31 0.31 1.5) shows that the magnetic excitation peak is at ~ 6 meV and that the background at low and high energy transfer is considerably lower than the neutron scattering at (0.31 0.31 1) between 4 and 10 meV. The solid lines are fits described in the text including the resolution function. The shaded areas are a guide to the eye.

The continuum scattering was investigated further using the triple-axis spectrometer DUALSPEC at Chalk River Laboratories and constant- $\vec{Q}$ scans were performed at 8.5 K for various wave-vectors in reciprocal space. The energy scans close to the AF point $l = 1$ revealed considerable continuum scattering above the well-defined Haldane excitation. The continuum extends up to 12 meV as shown in Fig. 2b) for the energy scan at (0.31 0.31 1). This wave-vector was chosen to be close to the 3D ordering wave-vector such that the Haldane excitation was at the lowest energy to give the largest possible energy window for the observation of the continuum. The scattering intensity at the AF zone boundary, $l = 1.5$, below and above the sharp excitation is considerably lower than the intensity at $l = 1$. The magnetic excitation was then fitted by a dispersion relation (an antisymmetrized Lorentzian weighted by the Bose factor) convoluted with the resolution ellipsoid given by Cooper-Nathans's expression [14]. The dispersion relation has been given by [4,6], and at 6 K the zone boundary energy is 6 meV, the

gap at $(0.81\ 0.81\ 1)$ is 1.23 meV [13] and the bandwidth of the gap along $[110]$ direction is given by the gap at $(0.33\ 0.33\ 1)$, which is 0.35 meV [15]. The Lorentzian half width of the excitation is 0.35 meV [13]. The fit gives an excellent account of the right-hand-side of the peak, as shown in Fig. 2b), but it cannot explain the slowly decreasing scattering continuum at energies above 4 meV.

The integrated intensity for the DUALSPEC data at the AF point $l = 1$ was inferred by scaling the peak intensity for $l = 1.5$ according to the l -dependence of the intensity measured by the MARI experiment. The integrated intensity of the continuum at 8.5 K is found to be about 12(3)% of the overall scattering at the AF wave-vector. This confirms the results obtained using the MARI spectrometer.

The temperature dependence of the continuum scattering was further investigated using the RITA triple-axis spectrometer at Risø and constant- $\vec{Q}$ scans were performed at the 1D point $(0.81\ 0.81\ 1)$. In Fig. 3 we show that the peak broadens and increases in energy with increasing temperature as reported elsewhere [13]. The well-defined peak was well fitted by an antisymmetrized Lorentzian weighted by the Bose factor and convoluted with the resolution ellipsoid. The flat background was determined from the scattered intensity below 25 K at the highest energy transfers measured ($E > 12$ meV), where the scattering was temperature independent and assumed to be non-magnetic. At low temperatures and energies higher than the well-defined peak energy, there was considerable continuum scattering (Fig. 3). After correction for neutron absorption, the total integrated intensity at 6 K and 12 K was about 14(3)% and 13(6)% of the integrated scattering at $l = 1$, respectively. At higher temperatures, the Lorentzian peak increases in energy and broadens so that the continuum could not be clearly identified. Nevertheless there was more scattering around 8 – 12 meV energy transfer at 6 and 9 K than at 30 K (see Fig. 3).

FIG. 3. Neutron scattering intensity at $l = 1$ wave-vector transfer as a function of energy transfer measured using the triple-axis spectrometer RITA in Risø for three different temperatures. The data is corrected for neutron absorption. The solid lines are the fits explained in the text including the spectrometer resolution and the shaded areas are a guide to the eye.

There are a number of theoretical predictions for the strength of a continuum at $l = 1$. A numerical diagonalization of a spin-1 chain with $N = 20$ sites and nearest-neighbor exchange interaction showed that at zero temperature the continuum above the Haldane excitation carries 3% of the total weight [16]. In the non-linear sigma model (NL σ M), a continuum arises from 3-particle scattering, and its integrated intensity is about 2% of the

spectral weight [17,18]. If the coupling of the magnetic chain is taken into account via a Random Phase Approximation (RPA), the 3-particle scattering becomes dependent on the wave-vector transfer perpendicular to the 1D axis, but its integrated intensity remains between 0.5 and 2.25% of the total spectral weight [18].

The portion of the intensity in the continuum scattering for CsNiCl₃ is about 12(2)% for $l = 1$. Our measurements for CsNiCl₃ show that the intensity between $l = 1$ and $l = 1.5$ reduces by a factor of 5.4, which is substantially lower than the theoretical prediction of 8.9. The absolute intensity of continuum scattering may therefore be only 7(2)% of the theoretical predicted intensity, but this is nevertheless much greater than the theoretical predictions.

A considerably stronger 3-particle continuum of about 17% is predicted by the Majorana Fermion Theory (MFT), which describes 1D magnetic systems in terms of a perturbation theory about the Heisenberg point of a model with biquadratic interactions as strong as the bilinear exchange [18]. It is unclear whether this model is applicable for CsNiCl₃. Furthermore, no indication of the pronounced peak at four times the gap energy, which is predicted for this model, was found in our measurements. Finally the MFT results are not in agreement with the numerical calculations of Takahashi [19].

In order to confirm that the continuum scattering arises from 3-particle scattering as predicted by the NL σ M we have searched for the onset of the continuum at 3Δ and for the pronounced maximum of the continuum at 6Δ as predicted by the NL σ M. With $\Delta = 0.94$ meV, our resolution is more than adequate to observe an increasing continuum intensity between $3\Delta = 2.8$ meV and $6\Delta = 5.6$ meV, but instead the spectrum shows a steadily decreasing intensity (Fig. 2, 3). This may arise because the 3D interactions in CsNiCl₃ cause a change in the excitation spectrum which is not properly taken into account via RPA and the pronounced maximum at 6Δ gets much broader as expected from kinematics. Possibly the effect can also account for the observed continuum intensity being higher than predicted for a 1D chain.

The observed continuum, Fig. 1, 2, 3, has intensity extending to approximately 12 meV, which is twice the maximum 1-particle energy for CsNiCl₃. Because similar measurements near $l = 0, 2$ show no evidence of scattering, we conclude that the continuum is strongest in a broad region where the AF fluctuations are largest. It is then possibly worth commenting that qualitatively the continuum scattering has features that are similar to the continuum found for $S=1/2$ systems (Fig. 1) [2].


It is clear that further theoretical work is needed to understand these results. It is unclear firstly whether a quantitative theory can be obtained from the 3-particle picture of the NL σ M or whether it is more appropriate to use another development such as the MFT approximation [21] to the Heisenberg model with biquadratic exchange

and secondly whether the form of the scattering depends on the inter-chain coupling or is characteristic of coupled chains.

In summary, we have measured the continuum scattering of the spin-1 chain compound CsNiCl_3 in a broad region that favours antiferromagnetic fluctuations and shown that it is stronger than expected for an ideal spin-1 chain both from numerical diagonalization and field theoretical calculations involving the $\text{NL}\sigma\text{M}$. These results are presented as observations from experiment, and while we cannot explain them we hope that they will stimulate further theoretical work.

We would like to thank Z. Tun for his assistance and Ian Affleck and A. M. Tsvelik for helpful discussions. Financial support for the experiments was provided by the EPSRC, by the EU through its Large Installations Program and by the British Council-National Research Council Canada Program. ORNL is managed for the U.S. D.O.E. by UT-Battelle, LLC, under contract no. DE-AC05-00OR22725. One of the authors (M. K.) is supported by a TMR-fellowship from the Swiss National Science Foundation under contract no. 83EU-053223.

-
- [1] F. D. M. Haldane, Phys. Rev. Lett. **50**, 1153 (1983).
 - [2] D. A. Tennant *et al.*, Phys. Rev. B **52**, 13368 (1995).
 - [3] G. Müller *et al.*, Phys. Rev. B **34**, 1429 (1981).
 - [4] W. J. L. Buyers *et al.*, Phys. Rev. Lett. **56**, 371 (1986).
 - [5] J. P. Renard *et al.*, Journal de Physique **49**, 1425 (1988).
 - [6] S. Ma *et al.*, Phys. Rev. Lett. **69**, 3571 (1992).
 - [7] F. D. M. Haldane, Phys. Lett. **93A**, 464 (1983).
 - [8] R. Morra *et al.*, Phys. Rev. B **38**, 543 (1988).
 - [9] A. A. Katori *et al.*, J. Phys. Soc. Jpn. **64**, 3038 (1995).
 - [10] M. Kenzelmann *et al.*, Quantum Spin Dynamics of Spin-1 Chain Compound CsNiCl_3 in its Paramagnetic Phase, to be published.
 - [11] I. Affleck and R. A. Weston, Phys. Rev. B **45**, 4667 (1992).
 - [12] D. Damle and S. Sachdev, Phys. Rev. B **57**, 8307 (1998).
 - [13] M. Kenzelmann *et al.*, Temperature Evolution of the Quantum Gap in CsNiCl_3 , to be published.
 - [14] M. J. Cooper and R. Nathans, Acta Cryst. **23**, 357 (1967).
 - [15] I. A. Zaliznyak *et al.*, Phys. Rev. B **50**, 15824 (1994).
 - [16] M. Takahashi, Phys. Rev. B **50**, 3045 (1994).
 - [17] M. D. P. Horton and I. Affleck, Phys. Rev. B **60**, 11891 (1999).
 - [18] F. H. L. Essler, cond-mat/9908186, 1999.
 - [19] M. Takahashi, Phys. Rev. Lett. **62**, 2313 (1989).
 - [20] M. Steiner *et al.*, J. Appl. Phys. **61**, 3953 (1987).
 - [21] A. M. Tsvelik, Phys. Rev. B **42**, 10499 (1990).



This figure "Fig3-three-scans-Risoe.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/cond-mat/0011046v1>

This figure "Fig2-MARI-C5-scans.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/cond-mat/0011046v1>